
Sirex noctilio: Overview of Biology and Control

Nathan Schiff
USDA Forest Service
Southern Research Station
Center for Bottomland Hardwoods Research
P.O. Box 227
Stoneville, MS 38776
nschiff@fs.fed.us







Normally Dennis Haugen would give this talk.

Dennis worked on *Sirex* noctilio for 8 years in Australia

He authored the Australian control plan

Also worked on *Sirex* noctilio in South America

He took many of the pictures used here.

I have studied the biology of native and exotic siricid woodwasps that attack hardwoods and conifers for the last 10 years.



I work with Dan Wilson a plant pathologist who studies wood decay fungi.




Sirex noctilio Classification: Hymenoptera: Symphyta, Siricoidea





Sirex noctilio



Host range:

Native Range: Europe, Asia, North Africa

Hosts: P. sylvestris, P. nigra, P. pinaster

Introduced Range:	Hosts
New Zealand 1900	P.radiata
Tasmania 1952	P.radiata
Australia 1961,	P.radiata
South Africa 1994,	P. radiata, P. patula
South America 1980-2001	
Brazil	P. taeda, P. elliottii
Argentina	P. banksiana, P. contorta
North America 2005	P. resinosa, P. strobus


taeda =loblolly, elliottii =slash, banksiana = Jack, resinosa =red, strobus = white, contorta = lodgepole, radiata =monterey, patula = mexican weeping, sylvestris = scots, nigra =austrian, pinaster= maritime



Biology:


Female *Sirex* mate with males in the crowns of trees and oviposit into the bole of stressed trees often half way up the tree.

Females also inject a wood decay fungus and a phytotoxic mucus into the wood.



Larva is "s" shaped with chewing mouthparts and postcornal spine.
Tunnels are packed with frass.

7-12 instars
Life cycle 1-2 years
Occasionally more than one generation per year
Larvae feed in sapwood and heartwood on fungus



Sirex noctilio indicators

Fungal Staining



Dripping Resin



Emergence Holes







Sirex noctilio kills North American pines. In Australian plantations up to 80% Mortality

In its native range *S. noctilio* is a minor pest like our siricids are here. In the southern hemisphere it is the most devastating pest of pines.

Will it be a problem in North America





Circumstances:

Range: Its in at least 20 counties in New York, 2 in Pennsylvania and in Ontario in the areas bounded by the Great Lakes (300 miles). (Speakers coming up will show delimiting maps)

Vulnerability: The larvae live within trees and would be very difficult to spray.

Dispersal: Adults are good flyers (maybe as much as 50 miles per year). Variable flight period so adults difficult to target.

Natural limits: No shortage of host material east of the plains.

We do not believe Sirex noctilio can be eradicated.

Clearly Sirex noctilio presents a threat to U.S. Forestry

What can we do about it?

What do the Southern Hemisphere countries do about Sirex noctilio?

The Australians developed a control program using:

Silvicultural control

Biological Control using

Parasitoid wasps

Rhyssine ichneumonids


Ibaliids

Parasitic nematode

Beddingia (=Deladenus) siricidicola



No thinning 68% loss, pre-commercial thinning no loss



Partial control with parasitoids


Megarhyssa sp.

Ibalia sp.

Parasitoids help but tend to be density independent.



NEMATODE LIFE CYCLE







Good control with nematode



Will the Nematode Work in North America? Ecological considerations:

Premise: The North American pine ecosystem is more ecologically complex than Southern Hemisphere pine plantations

Northern

100+ conifers in 12 genera 20 species native Siricidae Many conifer feeding insects Native wood decay fungi Native nematodes Many native parasitoids

Southern

Single pine monoculture
Single exotic siricid
Few pine feeding insects
One exotic fungus
One exotic nematode
Few introduced parasitoids

Some of the Possible Ecological Interactions

Tree x fungus:

Amylostereum wood decay studies

Fungus x fungus:

Somatic antagonism studies

Nematode x fungus:

Test nematodes on WDF

Nematode x insects:

Non target host-testing


Tree x Fungus interactions

How does *areolatum* compare to native *chailletii*36 species of conifers in 10 genera, 4 bottomland hardwoods

4 species of *Amylostereum* (3 isolates each) *areolatum, chailletii, laevigatum, ferreum*




Nematode x fungus: Test nematode survival and growth on WDF



Nematode x insects: Non target host-testing

Inoculate infested logs with fungus and nematodes. Dissect adults on emergence.



Monochamus titillator

Xylotrechus sagittatus

Will the Nematode Work in North America? Practical considerations:

Develop identification methods

Identify larvae with CO I

Microsatellites to locate source of *Sirex*

Microsatellites for fungal host race formation


Identify nematode species races??

Develop better traps and lures Chemistry, EAD's, Field testing

Develop trap tree technology

Best phenology for trap trees

Competitive exclusion (e.g. Sirex vs other insects)









Conclusions:

We have a new invasive exotic insect that kills pines.

It is likely to spread and can not be eradicated.

We have a classical biological control agent that has worked in Australia.

We have the plan the Australians used to control *Sirex noctilio*.

We need to do some research to adapt the system to North America.

It will be a long term program but it has the potential to be effective.

Acknowledgements:

USDA Forest Service FHP The State of New York **Animal Plant Health Inspection Service Oregon Department of Agriculture** Dan Wilson, Ted Leininger, Dennis Haugen, Don Duerr, Lynn Evans-Goldner, Jim Labonte Rick Hoebeke, Vic Mastro, David Smith, Bob Haack Ethan Angell, Yvonne DeMarino, Kevin Dodds, Sue Ellis, Harry Fulton, Paul Wrege, Noel Schneeburger Bob Mungari, Darryl Jewett, Jim Meeker, Brian Scheffler, Earl Taliercio, Edson Tadeu Iede, Dick Bashford, Meg Allen, Doug Street, Vicky Klasmer, Ron Billings and many others.

Special thanks to Amy Abel and Charisse Oberle